

# LAMPIRAN- LAMPIRAN

## Uji normalitas

		Tests of Normality					
		Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
	Jumlah penambahan	Statistic	df	Sig.	Statistic	df	Sig.
Rasa	X0 (30%)	.291	30	.000	.753	30	.000
	X1 (50)	.383	30	.000	.673	30	.000
	X2 (70%)	.227	30	.000	.862	30	.001
	X3 (90%)	.503	30	.000	.452	30	.000

a. Lilliefors Significance Correction

Anova satu arah

## ANOVA

Rasa

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	167.692	3	55.897	79.755	.000
Within Groups	81.300	116	.701		
Total	248.992	119			

Warna

		Tests of Normality					
		Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
	Jumlah penambahan	Statistic	df	Sig.	Statistic	df	Sig.
Warna	X0 (30%)	.357	30	.000	.729	30	.000
	X1 (50%)	.457	30	.000	.554	30	.000
	X2 (70%)	.488	30	.000	.492	30	.000
	X3 (90%)	.539	30	.000	.180	30	.000

a. Lilliefors Significance Correction

## Descriptives

Warna

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
X0 (30%)	30	4.03	.556	.102	3.83	4.24	3	5
X1 (50%)	30	3.73	.450	.082	3.57	3.90	3	4
X2 (70%)	30	3.20	.407	.074	3.05	3.35	3	4

X3 (90%)	30	1.97	.183	.033	1.90	2.03	1	2
Total	120	3.23	.896	.082	3.07	3.40	1	5

### ANOVA

Warna

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	74.867	3	24.956	140.526	.000
Within Groups	20.600	116	.178		
Total	95.467	119			

### Warna

Tukey HSD<sup>a</sup>

Jumlah penambahan	N	Subset for alpha = 0.05			
		1	2	3	4
X3 (90%)	30	1.97			
X2 (70%)	30		3.20		
X1 (50%)	30			3.73	
X0 (30%)	30				4.03
Sig.		1.000	1.000	1.000	1.000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 30.000.

Aroma

### Tests of Normality

	Jumlah penambahan	Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Aroma	X0 (30%)	.369	30	.000	.743	30	.000
	X1 (50%)	.344	30	.000	.755	30	.000
	X2 (70%)	.325	30	.000	.717	30	.000
	X3 (90%)	.537	30	.000	.275	30	.000

a. Lilliefors Significance Correction

### Descriptives

Aroma

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
X0 (30%)	30	3.80	1.157	.211	3.37	4.23	1	5
X1 (50%)	30	3.17	.592	.108	2.95	3.39	2	4
X2 (70%)	30	1.53	.571	.104	1.32	1.75	1	3
X3 (90%)	30	1.07	.254	.046	.97	1.16	1	2
Total	120	2.39	1.337	.122	2.15	2.63	1	5

### ANOVA

Aroma

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	152.292	3	50.764	97.655	.000
Within Groups	60.300	116	.520		
Total	212.592	119			

### Aroma

Tukey HSD<sup>a</sup>

Jumlah penambahan	N	Subset for alpha = 0.05		
		1	2	3
X3 (90%)	30	1.07		
X2 (70%)	30	1.53		
X1 (50%)	30		3.17	
X0 (30%)	30			3.80
Sig.		.064	1.000	1.000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 30.000.

Tekstur

### Tests of Normality

	Jumlah penambahan	Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Tekstur	X0 (30%)	.440	30	.000	.577	30	.000
	X1 (50%)	.423	30	.000	.597	30	.000
	X2 (70%)	.423	30	.000	.597	30	.000
	X3 (90%)	.354	30	.000	.637	30	.000

a. Lilliefors Significance Correction

### Descriptives

Tekstur

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
X0 (30%)	30	3.70	.466	.085	3.53	3.87	3	4
X1 (50%)	30	3.33	.479	.088	3.15	3.51	3	4
X2 (70%)	30	1.67	.479	.088	1.49	1.85	1	2
X3 (90%)	30	1.53	.507	.093	1.34	1.72	1	2
Total	120	2.56	1.083	.099	2.36	2.75	1	4

### ANOVA

Tekstur

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	112.492	3	37.497	160.505	.000
Within Groups	27.100	116	.234		
Total	139.592	119			

### Tekstur

Tukey HSD<sup>a</sup>

Jumlah penambahan	N	Subset for alpha = 0.05		
		1	2	3
X3 (90%)	30	1.53		
X2 (70%)	30	1.67		
X1 (50%)	30		3.33	
X0 (30%)	30			3.70
Sig.		.709	1.000	1.000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 30.000.

Tingkat Kesukaan

### Tests of Normality

Jumlah penambahan	Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.

Tingkat Kesukaan	X0 (30%)	.372	30	.000	.628	30	.000
	X1 (50%)	.409	30	.000	.599	30	.000
	X2 (70%)	.328	30	.000	.720	30	.000
	X3 (90%)	.412	30	.000	.648	30	.000

a. Lilliefors Significance Correction

### Descriptives

Tingkat Kesukaan

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
X0 (30%)	30	3.57	.679	.124	3.31	3.82	1	4
X1 (50%)	30	3.03	.556	.102	2.83	3.24	1	4
X2 (70%)	30	1.60	.563	.103	1.39	1.81	1	3
X3 (90%)	30	1.37	.556	.102	1.16	1.57	1	3
Total	120	2.39	1.102	.101	2.19	2.59	1	4

### ANOVA

Tingkat Kesukaan

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	104.092	3	34.697	99.380	.000
Within Groups	40.500	116	.349		
Total	144.592	119			

### Tingkat Kesukaan

Tukey HSD<sup>a</sup>

Jumlah penambahan	N	Subset for alpha = 0.05		
		1	2	3
X3 (90%)	30	1.37		
X2 (70%)	30	1.60		
X1 (50%)	30		3.03	
X0 (30%)	30			3.57

Sig.		.423	1.000	1.000
------	--	------	-------	-------

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 30.000.













