

## DAFTAR PUSTAKA

- Addakhil, A. R. (2021). *Formulasi dan evaluasi nanostructured lipid carrier ekstrak daun chrysanthemum cinerariifolium (trev.) vis* [Undergraduate, Universitas Islam Negeri Maulana Malik Ibrahim]. <http://etheses.uin-malang.ac.id/30094/>
- Adebisi, A. O., Kaialy, W., Hussain, T., Al-Hamidi, H., Nokhodchi, A., Conway, B. R., & Asare-Addo, K. (2016). Solid-state, triboelectrostatic and dissolution characteristics of spray-dried piroxicam-glucosamine solid dispersions. *Colloids and Surfaces B: Biointerfaces*, *146*, 841–851. <https://doi.org/10.1016/j.colsurfb.2016.07.032>
- Anggraeni, Y., Isnaeni, & Hendradi, E. (2017). Physical and Chemical Characteristics of Meloxicam from Nanostructured Lipid Carriers System Using Some Concentration Ratios of Monostearin and Alpha-Tocopherolacetate Lipid Matrix. *Asian Journal of Pharmaceutical and Clinical Research*, *10*(2), 132–137.
- Annisa, R., Hendradi, E., & Melani, D. (2016). Pengembangan sistem nanostructured lipid carriers (NLC) meloxicam dengan lipid monostearin dan miglyol 808 menggunakan metode emulsifikasi. *Journal of Tropical Pharmacy and Chemistry*, *3*(3), 156–169.
- Annisa, V. (2020). Review Artikel: Metode untuk Meningkatkan Absorpsi Obat Transdermal. *Journal of Islamic Pharmacy*, *5*(1), 18–27. <https://doi.org/10.18860/jip.v5i1.9157>
- Ansel, H. C., Allen Jr, L. V., & Popovich, N. G. (2020). *Ansel Bentuk Sediaan Farmasetis & Sistem Penghantaran Obat*.
- Baki, G., & Alexander, K. S. (2015). *Introduction to cosmetic formulation and technology*. John Wiley & Sons.
- Bhatia, M., & Rohilla, S. (2020). Formulation and optimization of quinoa starch nanoparticles: Quality by design approach for solubility enhancement of piroxicam. *Saudi Pharmaceutical Journal*, *28*(8), 927–935. <https://doi.org/10.1016/j.jsps.2020.06.013>

- Bindu, S., Mazumder, S., & Bandyopadhyay, U. (2020). Non-steroidal anti-inflammatory drugs (NSAIDs) and organ damage: A current perspective. *Biochemical Pharmacology*, *180*, 114147. <https://doi.org/10.1016/j.bcp.2020.114147>
- BPOM, R. (2014). Non-clinical toxicity test guidelines in in vivo (Indonesian: Pedoman Uji Toksisitas Nonklinik Secara In vivo). *Jakarta: Badan Pengawas Obat dan Makanan Republik Indonesia*.
- Bunjes, H., Westesen, K., & Koch, M. H. J. (1996). Crystallization tendency and polymorphic transitions in triglyceride nanoparticles. *International Journal of Pharmaceutics*, *129*(1), 159–173. [https://doi.org/10.1016/0378-5173\(95\)04286-5](https://doi.org/10.1016/0378-5173(95)04286-5)
- Chauhan, I., Yasir, M., Verma, M., & Singh, A. P. (2020). Nanostructured lipid carriers: A groundbreaking approach for transdermal drug delivery. *Advanced pharmaceutical bulletin*, *10*(2), 150.
- Chen-yu, G., Chun-fen, Y., Qi-lu, L., Qi, T., Yan-wei, X., Wei-na, L., & Guang-xi, Z. (2012). Development of a Quercetin-loaded nanostructured lipid carrier formulation for topical delivery. *International Journal of Pharmaceutics*, *430*(1), 292–298. <https://doi.org/10.1016/j.ijpharm.2012.03.042>
- Chu, D. H. (2012). Chapter 7. Development and Structure of Skin. Dalam L. A. Goldsmith, S. I. Katz, B. A. Gilchrest, A. S. Paller, D. J. Leffell, & K. Wolff (Ed.), *Fitzpatrick's Dermatology in General Medicine* (8 ed.). The McGraw-Hill Companies. [accessmedicine.mhmedical.com/content.aspx?aid=56021404](https://accessmedicine.mhmedical.com/content.aspx?aid=56021404)
- Cirri, M., Maestrini, L., Maestrelli, F., Mennini, N., Mura, P., Ghelardini, C., & Di Cesare Mannelli, L. (2018). Design, characterization and in vivo evaluation of nanostructured lipid carriers (NLC) as a new drug delivery system for hydrochlorothiazide oral administration in pediatric therapy. *Drug delivery*, *25*(1), 1910–1921.
- Ditjen POM. (2020) *Farmakope Indonesia*. Edisi Keenam. Jakarta: Departemen Kesehatan Republik Indonesia. Halaman 1403-1406
- Eroschenko, V.P. (2016). *Atlas Histologi*. Edisi Ke Dua Belas. Jakarta: Penerbit Buku Kedokteran EGC. Halaman 262-264

- Franz Cell—The Original*. (t.t.). PermeGear. Diambil 29 Januari 2022, dari <https://permegear.com/franz-cells/>
- Garcês, A., Amaral, M. H., Sousa Lobo, J. M., & Silva, A. C. (2018). Formulations based on solid lipid nanoparticles (SLN) and nanostructured lipid carriers (NLC) for cutaneous use: A review. *European Journal of Pharmaceutical Sciences*, *112*, 159–167. <https://doi.org/10.1016/j.ejps.2017.11.023>
- Gardouh, A. R., Faheim, S. H., Noah, A. T., & Ghorab, M. M. (2018). Influence of formulation factors on the size of nanostructured lipid carriers and nanoemulsions prepared by high shear homogenization. *International journal of pharmacy and pharmaceutical sciences*, 61–75.
- Ghadiri, M., Fatemi, S., Vatanara, A., Doroud, D., Najafabadi, A. R., Darabi, M., & Rahimi, A. A. (2012). Loading hydrophilic drug in solid lipid media as nanoparticles: Statistical modeling of entrapment efficiency and particle size. *International journal of pharmaceutics*, *424*(1–2), 128–137.
- Gowda, D., Tej, K., Moin, A., Anjali, Godugu, K., Patel, N., & Kamal, S. (2016). Nano structured lipid carrier based drug delivery system. *Journal of Chemical Research*, *8*, 627–643.
- Hariyadi, D. M., Rosita, N., & Rahayu, A. (2019). Design, optimization and characterization of glutathione loaded-alginate microspheres for topical antiaging. *Journal of Pharmacy & Pharmacognosy Research*, *7*(4), 223–233.
- Haryono, A., Restu, W. K., & Harmami, S. B. (2018). PREPARASI DAN KARAKTERISASI NANO PARTIKEL ALUMINIUM FOSFA. *Jurnal Sains Materi Indonesia*, *14*(1), 51–55.
- Hendradi, E., Purwanti, T., & Suryanto, A. A. (2012). Karakterisasi Sediaan Dan Uji Pelepasan Natrium Diklofenak Dengan Sistem Mikroemulsi Dalam Basis Gel HPC-M. *PharmaScientia I*.
- Hendradi, E., Rosita, N., & Rahmadhanniar, E. (2017). Effect of Lipid Ratio of Stearic Acid and Oleic Acid on Characteristics of Nanostructure Lipid Carrier (NLC) System of Diethylammonium Diclofenac. *Indonesian Journal of Pharmacy*, *28*(4), 198–204. <https://doi.org/10/C-03%20FULLTEXT.pdf>

- Hu, F.-Q., Jiang, S.-P., Du, Y.-Z., Yuan, H., Ye, Y.-Q., & Zeng, S. (2005). Preparation and characterization of stearic acid nanostructured lipid carriers by solvent diffusion method in an aqueous system. *Colloids and Surfaces B: Biointerfaces*, 45(3), 167–173. <https://doi.org/10.1016/j.colsurfb.2005.08.005>
- Katzung, B. G., Masters, S. B., & Trevor, A. J. (2014). Farmakologi Dasar & Klinik, Vol. 2, Edisi 12. *Editor Bahasa Indonesia Ricky Soeharsono et al., Penerbit Buku Kedokteran EGC, Jakarta.*
- Kovacevic, A., Savic, S., Vuleta, G., Müller, R. H., & Keck, C. M. (2011). Polyhydroxy surfactants for the formulation of lipid nanoparticles (SLN and NLC): Effects on size, physical stability and particle matrix structure. *International Journal of Pharmaceutics*, 406(1), 163–172. <https://doi.org/10.1016/j.ijpharm.2010.12.036>
- Lachman, L., Lieberman, H., & Kanig, J. L. (1994). Teori dan praktek Farmasi Industri II (Edisi 3). *Jakarta: Penerbit Universitas Indonesia.*
- Lai, F., Pini, E., Corrias, F., Perricci, J., Manconi, M., Fadda, A. M., & Sinico, C. (2014). Formulation strategy and evaluation of nanocrystal piroxicam orally disintegrating tablets manufacturing by freeze-drying. *International Journal of Pharmaceutics*, 467(1), 27–33. <https://doi.org/10.1016/j.ijpharm.2014.03.047>
- Lakshmi, P., & Kumar, G. A. (2010). Nanosuspension technology: A review. *Int J Pharm Sci*, 2(4), 35–40.
- Lambers, H., Piessens, S., Bloem, A., Pronk, H., & Finkel, P. (2006). Natural skin surface pH is on average below 5, which is beneficial for its resident flora. *International journal of cosmetic science*, 28(5), 359–370.
- Li, S., Su, Z., Sun, M., Xiao, Y., Cao, F., Huang, A., Li, H., Ping, Q., & Zhang, C. (2012). An arginine derivative contained nanostructure lipid carriers with pH-sensitive membranolytic capability for lysosomolytic anti-cancer drug delivery. *International journal of pharmaceutics*, 436(1–2), 248–257.
- Maciá Martínez, M.-Á. (2015). Economic Evaluation of the Restriction in the Use of Piroxicam in Spain. *Reumatología Clínica (English Edition)*, 11(6), 345–352. <https://doi.org/10.1016/j.reumae.2015.03.004>

- Maji, P., Gandhi, A., Jana, S., & Maji, N. (2013). Preparation and characterization of maleic anhydride cross-linked chitosan-polyvinyl alcohol hydrogel matrix transdermal patch. *J. PharmaSciTech*, 2(2), 62–67.
- Marwah, H., Garg, T., Goyal, A. K., & Rath, G. (2016). Permeation enhancer strategies in transdermal drug delivery. *Drug Delivery*, 23(2), 564–578. <https://doi.org/10.3109/10717544.2014.935532>
- Michael, P. (1995). *Ilmu Kimia*. Jakarta. Erlangga.
- Mohanraj, V. J., & Chen, Y. (2006). Nanoparticles-a review. *Tropical journal of pharmaceutical research*, 5(1), 561–573.
- Mu, L., & Feng, S. S. (2003). A novel controlled release formulation for the anticancer drug paclitaxel (Taxol®): PLGA nanoparticles containing vitamin E TPGS. *Journal of Controlled Release*, 86(1), 33–48. [https://doi.org/10.1016/S0168-3659\(02\)00320-6](https://doi.org/10.1016/S0168-3659(02)00320-6)
- Müller, R. H., Petersen, R. D., Hommoss, A., & Pardeike, J. (2007). Nanostructured lipid carriers (NLC) in cosmetic dermal products. *Advanced Drug Delivery Reviews*, 59(6), 522–530. <https://doi.org/10.1016/j.addr.2007.04.012>
- Naseri, N., Valizadeh, H., & Zakeri-Milani, P. (2015). Solid Lipid Nanoparticles and Nanostructured Lipid Carriers: Structure, Preparation and Application. *Advanced Pharmaceutical Bulletin*, 5(3), 305–313. <https://doi.org/10.15171/apb.2015.043>
- Navarro-Martínez, A., Vidal-Martínez, M., García-Rosa, I., Lázaro-Gómez, M. J., & Brotons-Román, J. (2015). Perfil de prescripción y adecuación del tratamiento con antiinflamatorios no esteroideos en el paciente diabético. *Revista de Calidad Asistencial*, 30(5), 256–264. <https://doi.org/10.1016/j.cali.2015.06.007>
- Negara, J. K., Sio, A. K., Rifkhan, R., Arifin, M., Oktaviana, A. Y., Wihansah, R. R. S., & Yusuf, M. (2016). Aspek mikrobiologis, serta Sensori (Rasa, Warna, Tekstur, Aroma) Pada Dua Bentuk Penyajian Keju yang Berbeda. *Jurnal Ilmu Produksi Dan Teknologi Hasil Peternakan*, 4(2), 286–290.
- Nuraeni, W., Daruwati, I., Widyasari, E. M., & Sriyani, M. E. (2013). *Verifikasi Kinerja Alat Particle Size Analyzer (PSA) Horiba LB-550 Untuk Penentuan Distribusi Ukuran Nanopartikel*.

- Penkina, A., Semjonov, K., Hakola, M., Vuorinen, S., Repo, T., Yliruusi, J., Aruväli, J., Kogermann, K., Veski, P., & Heinämäki, J. (2016). Towards improved solubility of poorly water-soluble drugs: Cryogenic co-grinding of piroxicam with carrier polymers. *Drug Development and Industrial Pharmacy*, 42(3), 378–388. <https://doi.org/10.3109/03639045.2015.1054400>
- Qian, C., Decker, E. A., Xiao, H., & McClements, D. J. (2012). Solid Lipid Nanoparticles: Effect of Carrier Oil and Emulsifier Type on Phase Behavior and Physical Stability. *Journal of the American Oil Chemists' Society*, 89(1), 17–28. <https://doi.org/10.1007/s11746-011-1882-0>
- Rahayu, A. (2020). Metode Validasi Flame Atomic Absorption Spectrometry (FAAS) Metode Destruksi Kering dan Destruksi Basah untuk Analisis Mineral Minuman Air Isotonik. *FARMASIS: Jurnal Sains Farmasi*, 1(1), 6–13.
- Rahayu, A., Rosyida, D. A. C., & Nuraini, I. (2022). FORMULASI DAN OPTIMASI NANOSTRUCTURED LIPID CARRIERS (NLC) KETOKONAZOL MENGGUNAKAN FULL FACTORIAL DESIGN: FORMULATION AND OPTIMIZATION OF NANOSTRUCTURED LIPID CARRIERS (NLC) KETOCONAZOLE USING FULL FACTORIAL DESIGN. *Medical Sains: Jurnal Ilmiah Kefarmasian*, 7(3), 561–570.
- Rahayu, A., Sari, D. P., & Ebtavanny, T. G. (2019). Design, Optimization and Characterization of Cefixime Microspheres. *Int J Pharma Res Heal Sci*, 7(5), 3051–3055.
- Rahmawan, T. G., Rosita, N., & Erawati, T. (2012). Characterisation of solid lipid nanoparticles p-methoxycinnamic acid (SLN-PMCA) formulated with different lipid component: Stearic acid and cetyl alcohol. *PharmaScientia*, 1(1), 23–29.
- Rohmah, M., Raharjo, S., Hidayat, C., & Martien, R. (2019). Formulasi dan Stabilitas Nanostructured Lipid Carrier dari Campuran Fraksi Stearin dan Olein Minyak Kelapa Sawit. *Jurnal Aplikasi Teknologi Pangan*, 8(1). <https://doi.org/10.17728/jatp.3722>

- Rowe, R. C., Sheskey, P., & Quinn, M. (2009). *Handbook of pharmaceutical excipients*. Libros Digitales-Pharmaceutical Press.
- Schäfer-Korting, M., Mehnert, W., & Korting, H.-C. (2007). Lipid nanoparticles for improved topical application of drugs for skin diseases. *Advanced Drug Delivery Reviews*, 59(6), 427–443. <https://doi.org/10.1016/j.addr.2007.04.006>
- Shah, M. R., Imran, M., & Ullah, S. (2017). *Lipid-based nanocarriers for drug delivery and diagnosis*. William Andrew.
- Shah, N. V., Seth, A. K., Balaraman, R., Aundhia, C. J., Maheshwari, R. A., & Parmar, G. R. (2016). Nanostructured lipid carriers for oral bioavailability enhancement of raloxifene: Design and in vivo study. *Journal of Advanced Research*, 7(3), 423–434. <https://doi.org/10.1016/j.jare.2016.03.002>
- Singhal, G., Patel, R., Prajapati, B., & Patel, N. (2011). Solid lipid nanoparticles and nano lipid carriers: As novel solid lipid based drug carrier. *International Research Journal of Pharmacy*, 2, 40–52.
- Sinko, P. J. (2011). Micromeritics. Dalam *Martin's Physical Pharmacy and Pharmaceutical Sciences: Physical Chemical and Biopharmaceutical Principles in the Pharmaceutical Sciences: Sixth Edition* (hlm. 442–468). Wolters Kluwer Health Adis (ESP).
- Sloane, E. (2004). *Anatomi dan Fisiologi untuk Pemula (diterjemahkan oleh Palupi Widyastuti)*. Jakarta.
- Souto, E. B., & Müller, R. H. (2007). Lipid nanoparticles (solid lipid nanoparticles and nanostructured lipid carriers) for cosmetic, dermal, and transdermal applications. Dalam *Nanoparticulate drug delivery systems* (hlm. 213–233). CRC Press.
- Thakker, K. D., & Chern, W. H. (2003). Development and validation of in vitro release tests for semisolid dosage forms-case study. *Dissolution Technologies*, 10(2), 10–16.
- Tjay, T. H., & Rahardja, K. (2007). *Obat-obat penting: Khasiat, penggunaan dan efek-efek sampingnya*. Elex Media Komputindo.
- Toll, R., Jacobi, U., Richter, H., Lademann, J., Schaefer, H., & Blume-Peytavi, U. (2004). Penetration Profile of Microspheres in Follicular Targeting of

- Terminal Hair Follicles. *Journal of Investigative Dermatology*, 123(1), 168–176. <https://doi.org/10.1111/j.0022-202X.2004.22717.x>
- Torchilin, V. P. (2006). *Nanoparticulates as drug carriers*. Imperial college press.
- Voigt, R. (1995). *Buku Pelajaran Teknologi Farmasi, Diterjemahkan oleh Soendani NS*. UGM Press, Yogyakarta.
- Yanhendri, S. W. Y. (2012). Berbagai bentuk sediaan topikal dalam dermatologi. *Cermin Dunia Kedokteran*, 194(6).
- Zhang, T., Chen, J., Zhang, Y., Shen, Q., & Pan, W. (2011). Characterization and evaluation of nanostructured lipid carrier as a vehicle for oral delivery of etoposide. *European journal of pharmaceutical sciences*, 43(3), 174–179.